



# Oligo - 8 Mn

Manganese Solution (14-0-0) + 8 Mn

COMPOSITION:	% w/w
Total Nitrogen(N)	14
Nitrogen from urea forms	14
Magnanese (Mn)	8

E.U. Origin

CONTENT: 1 Lt



■ **OLIGO-8Mn** is manganese solution for preventing/correcting manganese deficiency. Manganese is widely distributed in soils but in forms, which are not available to plants. The availability of Mn is strongly influenced by pH so that deficiency occurs in alkaline soils, sandy soils and in poor drained soils. Soil conditions, which favor Mn disorder, include:

- High organic matter.
- Imperfect drainage.
- High pH, especially following liming.
- Lack of consolidation.

■ **RATES FOR FOLIAR SPRAYS:**

Use 1-2 Lt / ton of water for foliar sprays, depending on plant growth stage.

■ **USERS:**

- **GRAPES:** Spray in spring when the shoots are 10 cm long. Repeat 2-3 times at 10 days intervals.
- **Citrus trees:** Spray 2-3 times at 10-15 days intervals, early in the spring when the new leaves expand. Avoid sprays close to blooming.
- **Fruit trees (Apples - Pears - Cherries- Plums):** Spray in spring, 2-3 times with the first symptoms, at 10-15 days intervals.
- **Vegetables (Tomato-Lettuce):** Apply when the first symptoms appear.
- **Lawn -Grasses:** Apply 2-3 times early in spring.
- **Corn-Oat-Barley:** Make 2-3 sprays, at 10-15 days intervals when some foliage has developed, (for the winter crops from March to bloom).
- **Beans-soybean-cowpeas:** Spray after the first 2 -3 leaves have expanded. Make 2 sprays at 15 days intervals. Applications are necessary when beans are growing on a sandy soil.
- **Potato-Sweet potato- tuber crops:** Potatoes may develop a mild Manganese deficiency. Spray 1-2 times when the first symptoms appear.

• **Sugar beet-Beets-root crops:** Spray 2-4 times at 10-15 days intervals, when the plants have obtained 3-4 leaves.

• **Cassava:** Spray with 1-2 Lt/ ton to supply plants with Manganese for an optimum growth.

• **Flowers-Ornamentals:** Spray with the first symptoms. Gerbera needs 1-2 preventing applications.

■ **COMPATIBILITY:** Compatible with most foliar fertilizers and agrochemicals. Do not use higher rates than those recommended. Avoid combinations with oily copper formulations.

■ **PRECAUTIONS:** Keep out of reach of children. Keep away of foodstuffs

■ **STORAGE:** Stored in room temperature the product remains stable for more than 3 years

Keep in original package.

DO NOT EXCEED THE RECOMMENDED RATES

Produced and Packaged by:  
**VIORYL S.A**  
Chemical and Agricultural Industry, Research S.A  
Athens, Greece  
Tel.: +30 22950 45100  
website: <http://www.vioryl.gr>